

Sedobassia sedoides (Pall.) Freitag & G. Kadereit in Slovakia: native species or alien weed?

Pavol Eliáš jun.^{1*}, Daniel Dítě²

¹Slovak University of Agriculture in Nitra, Slovakia

²Institute of Botany of the Slovak Academy of Sciences, Bratislava, Slovakia

The paper is aimed to explain occurrence and origin of *Sedobassia sedoides* in Slovakia because data presented in literature are not clear and complete. During our research, the occurrence of *Sedobassia sedoides* was confirmed in the Slovak flora on two reliable documented sites (Lednické Rovne, Bříňa) and one questionable location (Iliavka). Our research has confirmed that the species must be regarded as casual alien in the Slovak flora. Except cultivation in botanical garden of Lednické Rovne (West Slovakia, E: 18° 17' 13.04" N: 49° 17' 11.09") at the beginning of 19th century, the single occurrence was documented in 1872 in ruderal habitat of the Bříňa village. The distribution map is also presented.

Keywords: alien plants, distribution, flora, *Sedobassia sedoides*, Slovakia

1. Introduction

Bassia is a genus of flowering plants in the *Chenopodiaceae* family which are distributed in the western Mediterranean to eastern Asia including both annuals and perennial subshrubs (Aellen, 1979). The genus is defined also by its three types of C4 „kochioid“ leaf anatomy (Muhaidat et al., 2007). The *Bassia* species mainly occurs in steppe and desert ecosystems, some species can be found in ruderal sites and salt marshes up to subalpine altitudes (Collins and Blackwell, 1979; Mosyakin, 2004). In the Central Europe, three species are native (*Bassia sedoides*, *B. laniflora*, *B. prostrata*) and one species, *Bassia scoparia*, has spread here as invasive quarantine weed (Aellen, 1979; Jehlík, 1998).

Recently, the new genus *Sedobassia* Freitag & G. Kadereit was separated from traditionally accepted genus *Bassia* on the basis of different molecular sequences of ITS, rbcL, ATPB-rbcL and ndhF (Kadereit and Freitag, 2011). Those authors noted that isolated position of the genus *Sedobassia* in the molecular trees is corroborated by its peculiar C3/C4-intermediate leaf anatomy which is unique in *Camphorosmeae* (Carolin, 1983; H. Freitag, pers. obs.).

This new genus is monotypic – so it includes only single species *Sedobassia sedoides* (Pall.) Freitag and G. Kadereit [syn. *Kochia sedoides* Schrad., *Bassia sedoides* (Schrad.) Asch., Fig. 1]. The species is occurred from Hungary and countries of former Yugoslavia to Asian part of Russia, China and Mongolia (Zhu et al., 2003; Uotila, 2011). As an alien introduced also in the Czech Republic, Poland, Latvia, Estonia and Finland (Tomšovic, 1990; Sukhorukov




Figure 1 *Sedobassia sedoides*: A – fruit, B – habit of the upper part (Modified from Zhu et al., 2003)

*Correspondence: Pavol Eliáš jun., Slovak University of Agriculture in Nitra, Faculty of Agrobiobiology and Food Resources, Department of Botany, Tr. A. Hlinku 2, 949 76 Nitra, Slovakia, e-mail: pavol.elias.jun@gmail.com

and Uotila, 2007; Uotila, 2011) and it was reported from Algeria in the North Africa, too (Maire, 1962). The species was also reported from Slovakia (Dostál, 1991; Medvecká et al., 2012), but exact data have been still lacking. The paper is aimed to explain occurrence and origin of *Sedobassia sedoides* in Slovakia.

2. Material and methods

According to Zhu et al. (2003) *Sedobassia sedoides* is 10 – 60 cm tall annual herb; stem is erect, branched mostly from near middle, lanate-pilose. Leaves are linear, terete, 3 – 17 × 1 – 1.5 mm, fleshy, densely appressed pubescent and with a few villous hairs, base attenuate, apex obtuse. Flowers usually grouped in 2 or 3 per axillary glomerule. Perianth 5-lobed; abaxial appendages of segments triangular, subequaling the perianth. A fruit is smooth, broadly ovoid utricle.

The specimens of the species were studied in herbaria in Hungary (BP), the Czech Republic (BRNM, BRNU, MMI, MZ, OLM, PR, PRC) and in Slovakia (BRA, LTM, NI, KO, PMK, SAV, SLO and ZV). Herbarium abbreviations are according to Vozárová and Sutory (2001). Result of this study is presented on the point map. The map was designed by program ArcGis, version 9.2. The numbers of grids squares follow work of Niklfeld (1971). Coordinates of localities were taken from Google Earth. Phytogeographical divisions of Futák (1980) are also used. Nomenclature of flowering plants follows Marhold and Hindák (1998).

3. Results and discussion

During our study, only three data of *Sedobassia sedoides* were found in Slovakia (Fig. 2). Holuby (1888) mentioned it from the Lednické Rovne settlement in the Biele Karpaty Mts, but only as a cultivated plant. The species was cultivated there by A. Rochel, personal physician of the count of Johan Aspremont in years 1800 – 1811 and 1816 – 1820 (Vozárová, 2010). Rochel cultivated about 2,000 wild and ornamental plant species in botanical garden, among them also *Sedobassia sedoides*. This location represents the oldest occurrence of *S. sedoides* in Slovakia and it folded together with the destruction of the botanical garden in the mid – 19th century.

The second location was found in the Podunajská nížina Lowland. *Sedobassia sedoides* was collected here by L. Simonkai in September 1872 (Simonkai 1872 BP). The collector characterized the location as “in arenosis collium ad pagum Bény” = in sandy hill near the Bíňa village (Bény in Hungarian). Later Feichtinger (1899) stated the location of *B. sedoides* more exactly; it grew on the hill with the basilica in the centre of the village. The species has been occurring here only temporarily and later failed to confirm it (Domin, 1937; David, 2009). The presence of the species in this location has followed the occurrence in NW Hungary near the Esztergom settlement where some locations were known in the late 19th century. All those locations were not confirmed recently (Barina, 2003). Changes in land use are likely the reasons of termination of those locations, especially the




Figure 2 Distribution of *Sedobassia sedoides* in Slovakia in the past: ● – reliable documented locations in Lednické Rovne (1888) and Bíňa (1872), ○ – doubtful location in Ilávka (1989)

absence of grazing. Moderate and heavy grazing promotes spreading of annual xerophyllous plants of *Bassia sedoides* while extensive grazing or its absence caused the decline to extinction of the species at sites (Novikova et al., 2011).

The third location is doubtful. Dostál (1989) and Dostál and Červenka (1991) mentioned *S. sedoides* from the Iľávka village (now Iľavka, part of the Iľava town), but we found no reliable herbarium vouchered or literature reference to this data. Here are two possible explanations of this information: i) either he inaccurately transposed data of Holuby (1888) on the prevalence of *S. sedoides* near Lednické Rovne (this location is not so far from Iľavka) or ii) he found this species here but he published the finding not. It would be similar to the case of *Aegilops geniculata* which J. Dostál mentioned in his determination keys (Dostál, 1998; Dostál and Červenka, 1992), but herbarium voucher of this species was found in 2010 in the Dostál's unprocessed herbarium material (Eliáš et al., 2013).

If we want to evaluate the origin of *S. sedoides* in Slovakia, we must rely on data J. Dostál again. The author first presents an occurrence of the species in Slovakia as questionable (Dostál, 1948, 1954), but, despite lack of real data, he regarded it as native species of Slovakia in his later works (Dostál, 1989; Dostál and Červenka, 1991). Its occurrence was situated to saline habitats in the lower part of the Hron River (Dostál and Červenka, 1991), although the species was not mentioned in earlier or current works related to halophytic habitats in this area (e.g. Krist, 1940; Vicherek, 1973; Dítě et al., 2009). Inconsistent data on the origin of the species in our territory have led to it being introduced into alien species of Slovak flora (Medvecká et al., 2012). As the first occurrence the authors reported year 1948 – the year of the first issue of Dostál determination key to Czechoslovak flora (Dostál, 1948). We consider as a correct evaluation of the alien status of the species by Medvecká et al. (2012), viz. the inclusion of the species to the group of casual aliens of Slovakia. In addition to cultivation in the Botanic Garden in the Lednické Rovne village, *S. sedoides* is known only from location in the Biňa settlement. The character of the habitat – man-made raised mound with the Basilica, is clearly secondary and the species was introduced here as weed probably with draft animals of basilica visitors.

4. Conclusions

The occurrence of *Sedobassia sedoides* was confirmed in the Slovak flora on two reliable documented sites (Lednické Rovne, Biňa) and one questionable location (Iľavka). Our research has confirmed that the species was correctly included to casual weedy aliens of the Slovak flora, but the period of introduction is necessary to move more than 70 years to the past, to year 1872. Temporary re-introduction of *S. sedoides* in our territory can also not be excluded in the future.

5. Acknowledgement

We are thankful to the curators of herbaria visited during the study as well as to the anonymous reviewers for valuable comments to the manuscript. The research was made due the support of the Slovak Scientific Grant Agency VEGA (Grants No 2/0003/12 and 2/0008/13).

6. References

- AELLEN, P. (1979). *Chenopodiaceae*. In: REICHLINGER, K. H. (ed.) *Hegi G. Illustrierte Flora von Mitteleuropa III/2*. 2. ed. Berlin: Paul Parey, pp. 533–747.
- BARINA, Z. (2003) Details of Esztergom Danube floodplain flora. In *Kitaibelia*, vol. 8, no. 1, pp. 55–63 (in Hungarian).
- CAROLIN, R.C. (1983) The trichomes of the *Chenopodiaceae* and the *Amaranthaceae*. In *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie*, vol. 103, no. 4, pp. 451–466.
- COLLINS, S. L. and BLACKWELL, W. H. (1979) *Bassia* (*Chenopodiaceae*) in North America. In *Sida*, vol. 8, pp. 57–64.
- DAVID, S. (2009) The steppe flora and vegetation of the loess bulwarks in the Biňa village (district Nové Zámky, SW Slovakia). In *Bulletin Slovenskej botanickej spoločnosti*, vol. 31, no 2, pp. 77–85 (in Czech).
- DÍTĚ, D., ELIÁŠ, P. jun. and ŠUVADA, R. (2009) The current distribution and status of community *Puccinellietum limosae* in Slovakia. In *Thaiszia – Journal of Botany*, vol. 19, no. 1–2, pp. 63–70.
- DOMIN, K. (1937) Vegetation of „Avar Hrink“ near Biňa in the southern Slovakia. In *Věda Přírodní*, vol. 18, pp. 164–166 (in Czech).
- DOSTÁL, J. (1948) *Flora of Czechoslovakia and illustrated key to the identification of all vascular plants in Czechoslovakia wild or commercially cultivated*. Praha: Nakladatelství ČSAV (in Czech).
- DOSTÁL, J. (1954) *Key to Complete Flora Czechoslovakia*. Praha: Nakladatelství ČSAV (in Czech).
- DOSTÁL, J. (1989) *New flora of Czechoslovakia I*. Praha: Academia (in Czech).
- DOSTÁL, J. and ČERVENKA, M. (1991) *Big key for determining of higher plants I*. Bratislava: SPN (in Slovak).
- DOSTÁL, J. and ČERVENKA, M. (1992) *Big key for determining higher plants II*. Bratislava: SPN (in Slovak).
- ELIÁŠ, P. jun. et al. (2013) Distribution and origin of *Aegilops* species in Slovakia. In *Thaiszia – Journal of Botany*, vol. 23, no. 2, p. 117–129.
- FEICHTINGER, S. (1899) *The Esztergom county and the surrounding flora*. Esztergom: Esztergomvidéki Régészeti és Történelmi Társaságkiadványa (in Hungarian).
- FUTÁK, J. (1980) Phytogeographical division of Slovak Socialist Republic (1 : 1 000 000). In: MAZÚR, E. (ed.) *Atlas of the Slovak Socialist Republic*. Bratislava: SAV, p. 88 (in Slovak).
- HOLUBY, J. L. (1888) *Flora des Trencsiner Comitatus*. Trencsin.
- JEHLÍK, V. (ed.) (1998) *Alien expansive weeds of the Czech Republic and the Slovak Republic*. Praha: Academia.
- KADEREIT, G. and FREITAG, H. (2011) Molecular phylogeny of *Camphorosmeae* (*Camphorosmoideae*, *Chenopodiaceae*): Implications for biogeography, evolution of C4-photosynthesis and taxonomy. In *Taxon*, vol. 60, no. 1, pp. 51–78.

- KRIST, V. (1940) Halophytic vegetation of SW Slovakia and the northern part of the Small Hungarian Plain. In *Práce moravské přírodovědecké společnosti*, vol. 12, no. 10, pp. 1–100 (in Czech).
- MAIRE, R. (1962) *The flora of North Africa*. Paris: Lechevalier éd.
- MARHOLD, K. and HINDÁK, F. (eds.) (1998) *Checklist of non-vascular and vascular plants of Slovakia*. Bratislava: Veda.
- MEDVECKÁ, J. et al. (2012) Inventory of the alien flora of Slovakia. In *Preslia*, vol. 84, no. 2, pp. 257–309.
- MOSYAKIN, S. (2004) *Bassia* Allioni. In: Flora of North America Editorial Committee (eds.) *Flora of North America* Vol. 4. New York and Oxford, pp. 309–310.
- MUHAI DAT, R., SAGE, R.F. and DENGLER, N.G. (2007) Diversity of Kranz anatomy and biochemistry in C4 eudicots. In *American Journal of Botany*, vol. 94, no. 3, pp. 362–381. DOI: 10.3732/ajb.94.3.362.
- NIKL FELD, H. (1971) Bericht über die Kartierung der Flora Mitteleuropas. In *Taxon*, vol. 20, pp. 545–571.
- NOVIKOVA, N. M. et al. (2011) Changes of solonetz complex plant cover under different stocking pressure at Dzhanybek scientific station and adjacent territories. In *Arid Ecosystems*, vol. 1, no. 4, pp. 290–298. DOI: 10.1134/S2079096111040123.
- SUKHORUKOV, A. P. and UOTILA, P. (2007) Additions and corrections to the alien Chenopodiaceae flora of Finland and NW Russia. In *Memoranda Societatis pro Fauna et Flora Fennica*, vol. 83, no. 2, pp. 53–58.
- TOMŠOVIC, P. (1990) *Bassia* Allioni. In: HEJNÝ, S. and SLAVÍK, B. (eds.) *Flora of the Czech Republic 2*. Praha: Academia, pp. 285 (in Czech).
- UOTILA, P. (2011) *Chenopodiaceae* (pro parte majore). Euro+Med Plantbase – the information resource for Euro-Mediterranean plant diversity. [Online] Available from: <http://www.emplantbase.org/home.html/>. [Accessed: 2013-11-25].
- VICHEREK, J. (1973) Die Pflanzengesellschaften der halophyten und subhalophyten Vegetation der Tschechoslowakei. In *Vegetace ČSSR*, vol. A, no. 5, pp. 1–200.
- Vozárová, M. (2010). Rochel, Anton. In: Vozárová, M. and Šípošová, H. (eds.) *Personalities of Botany in Slovakia*. Bratislava: Veda, pp. 439–441 (in Slovak).
- VOZÁROVÁ, M. and SUTORÝ, K. (2001) Index of herbaria in the Czech Republic and the Slovak Republic. In *Bulletin Slovenskej botanickej spoločnosti*, Suppl. 7, pp. 1–95 (in Czech and Slovak).
- ZHU, G., MOSYAKIN, S. L. and CLEMANTS, S. E. (2003) *Bassia* Allioni. In: WU, Z. Y., RAVEN P. H. and HONG, D. Y. (eds.) *Flora of China*. Vol. 5. (*Ulmaceae through Basellaceae*). Beijing and St. Louis: Science Press and Missouri Botanical Garden Press, pp. 386–387.